



and outlined with warm buff and velvety fuscous black, which extends to postmedial fascia as a short dash; claviform mark outlined in fuscous black; antemedial fascia wavy, warm buff edged with fuscous black; postmedial, excurved, crenulate line of fuscous black and warm buff, commencing with a small, warm, buff patch at costa; subterminal fascia irregular, interrupted, warm buff preceded by a series of indistinct, interneural, fuscous, sagittate marks (points basad); a broad, ill-defined fuscous bar from postmedial nearly to termen, on, and on each side of, vein 5, but not touching 4 or 6; a terminal series of velvety, fuscous black lunules; fringe pecan brown, warm buff at base succeeded by a zone of neutral red. Underside: fuscous bordered with neutral red, postmedial and subterminal bands indicated, pale along inner margin and with a streak of warm buff below costa at basal third. *Hindwing*: light buff suffused with fuscous, veins fuscous, and area beyond postmedial more strongly suffused with fuscous; underside, lower half light buff, upper half suffused with neutral red, subterminal band fuscous, postmedial fuscous, strongly indicated from costa to vein 6, a diffuse fuscous discocellular lunule.

*Expanse* 44mm. (Tip to tip 42mm.)

*Holotype*. *Male*. Haight's Place, Pauai, subprov. Benguet, Luzon, I.,  
7,000 ft. 8.XI.1912.

Nearest ally *M. pannosa*, Moore (India, Ceylon).

### **Cirphis lipara**, sp.n.

*Male*.—Palpus: light buff, tinged with snuff brown above. Antenna: ciliated, with paired setae, light buff on basal quarter. Head: frons light buff, fuscous laterally, vertex light buff, tinged with snuff brown. Thorax: light buff, patagium light buff tinged with snuff brown, with two narrow transverse bands of fuscous black, snuff brown posteriorly, tegula light buff tinged with snuff brown, with a few fuscous black scales. Abdomen; light buff, with basal dorsal crest tinged with snuff brown, venter light buff, with fuscous black patch wide at base, diminishing to a point posteriorly, and a fuscous black spot on sides of each of segments 6 to 8, anal tuft light buff. *Forewing*: snuff brown, with cartridge buff band on and below costa, and veins 6 to 12; white on median nervure at distal half of cell, continuing along veins 3 and 4; area below cell up to postmedial fascia cartridge buff tinged with snuff brown (except just at wing-base), sparsely irrorated with a number of scattered fuscous black scales; a white and cartridge buff, diagonal, wedge-shaped fascia from apex, widest at inner margin between medial and postmedial fasciae, its outer edge overlapping the postmedial series of fuscous black, interneural points from vein 5 to anal vein; area between this line and termen snuff brown; interneural fuscous black spots on termen. Underside: cartridge buff overlaid with opalescent scales; upper half suffused with pinkish vinaceous, a prominent fuscous black spot on costa between end of cell and wing-apex, a trace of the diagonal wedge-shaped fascia, and a terminal series of interneural fuscous black points. *Hindwing*: upper side cartridge buff suffused with fuscous between veins 2 and 6; underside cartridge buff overlaid with opalescent scales, with a few fuscous scales on discocellulars and between them and wing-apex.

*Expanse* 36mm. (Tip to tip 35mm.)

<i>Holotype. Male.</i>	Haight's Place, Panai, subprov. Benguet, Luzon, I., 7000 ft.	30.XI.1912.
<i>Paratype. Male.</i>	" " "	" " " 26.VII.1913.
<i>Paratype. Male.</i>	" " "	" " " 28.VII.1913.
<i>Paratype. Male.</i>	Bagnio, subprov. Benguet, Luzon.	5000 ft. 1.IV.1913.

This species is similar to *C. decissima*, Wlk., but differs in having the diagonal line from apex more sharply defined, and the white scaling on median nervure and veins 3 and 4 is more extensive.

### *Cirphis albomarginalis*, sp.n.

*Male*.—Palpus: cartridge buff, with fuscous above. Antenna: ciliated, with paired setae. Head: frons and vertex cartridge buff and fuscous mixed. Thorax: cartridge buff, a tuft of pale drab grey speckled with fuscous hair-scales on mesothorax; patagium, cartridge buff anteriorly, edged with fuscous, cartridge buff tinged with wood brown posteriorly, tegula cartridge buff tinged with wood brown with a few fuscous black scales intermixed. Abdomen: drab with prominent fuscous crest on basal segment, other crests cartridge buff; venter cartridge buff, with a row of three half hidden fuscous spots at sides; anal-tuft cartridge buff. Pectus: cartridge buff. Legs: foreleg cartridge buff with fuscous inwardly, other legs cartridge buff speckled with fuscous, fuscous on spurs. *Forewing*: wood brown, cartridge buff band along and below costa, with scattered fuscous scales and three fuscous spots on costa, sub-basally, antemedially and postmedially; veins 6 to 12 cartridge buff, distal two thirds of cell cartridge buff lightly irrorated with wood brown, orbicular faintly marked by a wood brown spot, reniform ill-defined by fuscous; median nervure white, the white continuing along veins 3 and 4; a diagonal fascia from apex to inner margin at two thirds, from apex to vein 5, broad, cartridge buff, continuing as a broken row of postmedial fuscous dots; interneural, fuscous spots on termen; cartridge buff patch near tornus; underside cartridge buff, with scattered fuscous scales below costa, fuscous mark on discocellulars, a broken, fuscous, postmedial band with a fuscous spot at costa, a fuscous postmedial to subterminal patch between veins 2 and 5. *Hindwing*: fuscous, cartridge along costa and cilia; underside cartridge buff sparsely irrorated with fuscous; fuscous mark on discocellulars, fuscous dots on veins postmedially, interneural fuscous dots terminally.

(*Expanse* 38mm. (Tip to tip 36mm.))

*Female*.—Similar to male, slightly paler.

(*Expanse* 38mm. (Tip to tip 36mm.))

<i>Holotype. Male.</i>	Haight's Place, Panai, subprov. Benguet, Luzon, I., 7,000 ft.	12.V.1913.
<i>Allotype. Female.</i>	" " "	" " " 7,000 ft. 13.I.1912.
<i>Paratype. Female.</i>	Bagnio, subprov. Benguet, Luzon.	5,000 ft. 20.IV.1912.
<i>Paratype. Female.</i>	" " "	" " " 1.V.1913.

